

# Instantons and the ADHM Construction

## Lecture 1

Alex Atanasov

December 6, 2016

### Abstract

We explore connections on  $\mathbb{R}^4$  and the Yang-Mills equations arising from minimizing a quantity known as action. We study solutions to these equations possessing nonzero action, known as instantons, and demonstrate a method to construct all instantons on  $\mathbb{R}^4$  with dimension  $n$  and topological charge  $k$ . This is the ADHM construction of Atiyah et al.

## 1 Motivation

In this course we have seen examples of geometrization: the association of geometric structure to an underlying algebraic structure. We've seen that categorification of  $\mathfrak{sl}_q(2, \mathbb{C})$  gives rise to cohomology rings of Grassmanians. In a similar vein, more general affine Lie algebras  $\hat{\mathfrak{g}}$  give rise to geometric spaces that can be understood as moduli spaces of instantons on asymptotically-locally-euclidean (ALE) spaces  $\mathbb{C}^2/\Gamma$ , in one-to-one correspondence with the extended Affine Dynkin diagrams.

We give an introduction to instanton construction first in the simple case of  $\mathbb{C}^2 \cong \mathbb{R}^4$ . Even in this simple case, we will see how this theory is deeply connected to affine Lie algebras, Hilbert schemes, and quiver varieties.

## 2 Yang Mills Instantons on $\mathbb{R}^4$

### 2.1 Connection and Curvature Forms

**Definition 2.1.** A **Hermitian vector bundle**  $\pi : E \rightarrow M$  over a base space  $M$  is a complex vector bundle over  $M$  equipped with a Hermitian inner product on each fiber.

Yang Mills theory on  $M$  concerns itself with the metric-compatible **connections**  $A$  on  $E$ .

**Definition 2.2** (Connection on a Vector Bundle). A connection  $A$  on a vector bundle  $\pi : E \rightarrow M$  of rank  $n$  is a  $\mathfrak{gl}(n)$ -valued 1-form

For a Hermitian bundle, we restrict to  $\mathfrak{u}(n)$ , to work with only metric-compatible connections. Each such connection  $A \in \mathcal{A}$  is a  $\mathfrak{u}(n)$ -valued 1-form acting on  $E$  by  $\rho$ .

**Definition 2.3** (Covariant Exterior Derivative). For a given connection  $A \in \Omega^1(M, \mathfrak{u}(n))$ , we obtain a corresponding differential operator on  $M$ :

$$d_A := d + \rho(A) \tag{1}$$

**Observation 2.4.** *In coordinate language, we can write:*

$$(d_A)_\mu = \partial_\mu + \rho(A_\mu) \tag{2}$$

We can then define the **curvature** 2-form by having this derivative act on its own connection 1-form

**Definition 2.5** (Curvature/Field-Strength 2-form).

$$\begin{aligned} F &:= d_A A = dA + A \wedge A \\ &= dA + \frac{1}{2}[A, A] \end{aligned} \tag{3}$$

**Observation 2.6.** *In coordinate language, we can write:*

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu] \tag{4}$$

$$s.t. F = \frac{1}{2} F_{\mu\nu} dx^\mu \wedge dx^\nu \tag{5}$$

We conclude with an identity that can be checked by direct computation

**Proposition 2.7** (Bianchi Identity).

$$d_A F = 0 \tag{6}$$

## 2.2 The Action

For our purposes,  $M = \mathbb{R}^4$  will be the manifold in question. In particular  $\mathbb{R}^4$  has Riemannian structure, so we are given the Hodge-star operator

$$\star : \Omega^k \rightarrow \Omega^{n-k}.$$

We define the **action**, from which we will obtain all information about the dynamics, by:

$$S_E[\mathcal{A}] = - \int_M \text{Tr}(F \wedge \star F) \quad (7)$$

**Proposition 2.8.**  $\text{Tr}(F \wedge \star F)$  is globally-defined and gauge invariant

*Proof.* This follows directly from the cyclic properties of the trace, and the transformation laws on  $F$  making it transform under the adjoint representation.  $\square$

We want to find  $A$  so that  $S_E[\mathcal{A}]$  is a minimum. To do this, we use standard calculus of variations. Consider a local perturbation  $A + t\alpha$

$$\begin{aligned} F[A + t\alpha] &= d(A + t\alpha) + A \wedge A + t[A, \alpha] + O(t^2) \\ &= F[A] + t(d\alpha + [A, \alpha]) + O(t^2) \\ &= F[A] + d_A\alpha + O(t^2) \end{aligned} \quad (8)$$

so that to order  $t$ :

$$\begin{aligned} \|F[A + t\alpha]\|^2 &= \|F[A + t\alpha]\|^2 + 2t(F[A], d_A\alpha) \\ &\Rightarrow (F[A], d_A\alpha) = 0 \quad \forall \alpha \end{aligned} \quad (9)$$

By taking adjoints, this gives:

$$\begin{aligned} &\Rightarrow \star d_A \star F[A] = 0 \\ &\Rightarrow d_A \star F = 0 \end{aligned} \quad (10)$$

This, together with the tautological Bianchi identity:  $d_A F = 0$  form the Yang-Mills equations. These equations are very difficult to solve in all but abelian gauges, where they become linear.

## 2.3 Instantons and Topological Charge

**Proposition 2.9.** Let  $\dim M = 4$ . Then  $\int_M \text{Tr}(F \wedge F)$  is independent of changes in  $A$ .

*Proof.* Following the same variational procedure will give us  $d_A F$ , which is zero always, independent of any condition on  $A$ .  $\square$

We define the **topological charge**  $k$  of the theory by

$$k := -\frac{1}{8\pi^2} \int_M \text{Tr}(F \wedge F) \quad (11)$$

**Proposition 2.10.** When  $M = S^4$ , we have that  $k$  is an integer.

*Proof.* The proof lies in simple ideas from Chern classes and classifying bundles over  $S^4$ . It establishes a one-to-one correspondence between the global topology type of the bundle  $E$  over  $S^4$  and the topological charge. □

Now note that on  $\mathbb{R}^4$ , we have  $\star\star = 1$ . This means that  $\star$  has eigenvalues  $\pm 1$  and so  $\Omega^2(U, \mathfrak{g})$  splits as a direct sum of two orthogonal spaces:

$$\Omega^2(\mathbb{R}^2, \mathfrak{u}(n)) = \Omega_+^2 \oplus \Omega_-^2 \quad (12)$$

called **self-dual** and **anti-self-dual** spaces respectively.

We can “symmetrize” any form to become a sum of a self-dual and an anti-self dual one. In particular, if we write:

$$F = F_+ + F_- \quad (13)$$

then we have

$$\begin{aligned} -8\pi^2 k &= \int_M \text{Tr}[(F_+ + F_-) \wedge (F_+ + F_-)] d\text{Vol} \\ &= \int_M \text{Tr}[(F_+) \wedge (F_+)] d\text{Vol} + \int_M \text{Tr}[(F_-) \wedge (F_-)] d\text{Vol} \\ &= \int_M \|F_+\|^2 d\text{Vol} - \int_M \|F_-\|^2 d\text{Vol} \end{aligned} \quad (14)$$

Note that the absolute value of this gives:

$$8\pi^2 k \leq \int_M \|F\|^2 = |S_A[F]| \quad (15)$$

**Proposition 2.11.** *The action is bounded below by this topological charge and is in fact equal to it exactly when one of  $F_+ = 0$  or  $F_- = 0$ .*

We call a solution an **instanton** of the theory. Its action is equal to the topological charge, and in fact we call this the **instanton number** when appropriate. We are interested in the space of instantons modulo gauge equivalence

**Definition 2.12.** The **gauge group**  $\mathcal{G}$  of all metric-compatible transformation on  $E$ , restricts to  $\text{SU}(n)$  at each point. Two connections  $A_1, A_2$  are Gauge equivalent if they differ by an element in  $\mathcal{G}$ . We are interested in the space of connections modulo gauge.

Instantons on  $\mathbb{R}^4$  must have that  $F$  is either self-dual or anti-self-dual. In the latter case:

$$\star F = -\star F \quad (16)$$

This equation is much simpler to solve than the equation of motion  $d_A \star F = 0$ . The anti-self-duality (ASD) equations can be written out explicitly:

$$\begin{aligned} F_{12} + F_{34} &= 0 \\ F_{14} + F_{23} &= 0 \\ F_{13} + F_{42} &= 0 \end{aligned} \quad (17)$$

This can also be written in terms of commutators of the covariant derivatives. If we denote  $(d_A)_\mu$  simply by  $D_\mu$  then  $F_{\mu\nu} = (d_A)_\mu(d_A)_\nu = [D_\mu, D_\nu]$ .

$$\begin{aligned} [D_1, D_2] + [D_3, D_4] &= 0 \\ [D_1, D_4] + [D_2, D_3] &= 0 \\ [D_1, D_3] + [D_4, D_2] &= 0 \end{aligned} \tag{18}$$

**Proposition 2.13.** *There are no instantons on Minkowski space  $\mathbb{R}^{3,1}$ .*

*Proof.*  $\star\star = -1$  on Minkowski space, so  $\star$  has eigenvalues  $\pm i$ , meaning the duality equations would require  $\star F = \pm iF$ , but  $F \in \Omega^2(\mathbb{R}^4, \mathfrak{u}(n))$  is a real object.  $\square$

**Proposition 2.14.** *For all connections on a given vector bundle  $E$ , the instanton number is an invariant.*

*Proof.* This follows since for instantons  $S_A = 8\pi k$  is independent of the connection.  $\square$

**Corollary 2.15.** *There are no instantons when  $G$  is abelian.*

*Proof.*  $F = dA \Rightarrow \|F\| = (\star dA, dA) = (\delta \star A, dA) = (\star A, d^2 A) = 0$   $\square$

We then have two invariants to note:  $n$  and  $k$ . We will be especially interested in the moduli space of all instantons for specific  $n$  and  $k$  (modulo gauge). From now on, we will focus specifically on anti-self-dual (ASD) instantons.

$$\mathcal{M}_{ASD}(n, k)$$

Self-dual instantons can be constructed in a straightforward one-to-one manner from the ASD instantons.

There is one subtlety: For  $k$  to be finite, we need  $F$  to vanish sufficiently quickly. This gives a bound for  $|F| = |d_A A(x)| = O(|x|^{-4})$  for large  $x$ . This further gives a constraint on the gauge group  $\mathcal{G}$  as  $x \rightarrow \infty$  to have locally trivial structure. Instantons with this condition on their behaviour and gauge group are called **framed** instantons.

We say that in a neighborhood of infinity of  $S^4$ , the gauge group element must give a section of the bundle  $E$  that has a local trivialization  $\Phi : E_\infty \rightarrow \mathbb{C}^n$ . We denote the moduli space of framed instantons by

$$\mathcal{M}_{ASD}^{fr}(n, k)$$

### 3 The ADHM Construction

#### 3.1 The Data

Let  $x_1, x_2, x_3, x_4$  parameterize a  $\mathbb{R}^4$ , and write this as  $\mathbb{C}^2$  using  $z_1 = x_2 + ix_1, z_2 = x_4 + ix_3$ . We can then write all the  $(d_{\mathcal{A}})_{\mu}$  (from now on just  $D_{\mu}$ ). Moreover in terms of the complex coordinates, we get

$$\begin{aligned} \mathcal{D}_1 &= \frac{1}{2}(D_2 - iD_1) \\ \mathcal{D}_2 &= \frac{1}{2}(D_4 - iD_3) \end{aligned} \tag{19}$$

We can express anti-self duality of  $F_{\mu\nu}$  in terms of these  $\mathcal{D}_{\mu}$  through two equations:


$$\begin{aligned} [\mathcal{D}_1, \mathcal{D}_2] &= 0 \\ [\mathcal{D}_1, \mathcal{D}_1^{\dagger}] + [\mathcal{D}_2, \mathcal{D}_2^{\dagger}] &= 0 \end{aligned} \tag{20}$$

The idea behind ADHM is to convert these  $D_i$  to matrices  $B_i$  in a method akin to taking “Fourier transforms”, and adding source terms depending on  $k$ .

**Definition 3.1** (ADHM Data). Let  $U$  be a 4-dimensional space with complex structure. An **ADHM System** on  $U$  is a set of linear data:

1. Vector spaces  $V, W$  over  $\mathbb{C}$  of dimensions  $k, n$  respectively.
2. Complex  $k \times k$  matrices  $B_1, B_2$ , a  $k \times n$  matrix  $I$ , and an  $n \times k$  matrix  $J$ .

We can see this diagrammatically by the following doubled, framed quiver:



**Definition 3.2** (ADHM System). A set of ADHM Data is an ADHM system if it satisfies the following constraints:

1. The ADHM equations:

$$\begin{aligned} [B_1, B_2] + IJ &= 0 \\ [B_1, B_1^{\dagger}] + [B_2, B_2^{\dagger}] + II^{\dagger} - J^{\dagger}J &= 0 \end{aligned} \tag{21}$$

These quantities are called real and complex moment maps, respectively.

2. For any two  $x, y \in \mathbb{C}^2$  with  $x = (z_1, z_2), y = (w_1, w_2)$ , the map:

$$\alpha_{x,y} = \begin{pmatrix} w_2 J - w_1 I^\dagger \\ -w_2 B_1 - w_1 B_2^\dagger - z_1 \\ w_2 B_2 - w_1 B_1^\dagger + z_2 \end{pmatrix} \quad (22)$$

is injective from  $V$  to  $W \oplus (V \otimes U)$  while

$$\beta_{x,y} = \begin{pmatrix} w_2 I + w_1 J^\dagger & w_2 B_2 - w_1 B_1^\dagger + z_2 & w_2 B_1 + w_1 B_2^\dagger + z_1 \end{pmatrix} \quad (23)$$

is surjective from  $W \oplus (V \otimes \mathbb{C}^2)$  to  $V$ .

It's worth noting that  $W \oplus (V \otimes \mathbb{C}^2) \cong W \oplus V \oplus V$ .

**Lemma 3.3.** *If  $(B_1, B_2, I, J)$  satisfy the above conditions, then for  $g \in U(k)$ , we get*

$$(gB_1g^{-1}, gB_2g^{-1}, gI, Jg^{-1}) \quad (24)$$

also satisfy the above conditions.

Thus we care about solutions to these equations modulo  $U(V)$ .

*Proof.* It's a quick check through direct algebra that the equations are preserved.  $\square$

**Proposition 3.4.** *The ADHM equations are satisfied iff*

$$V \xrightarrow{\alpha_{x,y}} W \oplus (V \otimes \mathbb{C}^2) \xrightarrow{\beta_{x,y}} V \quad (25)$$

is a complex

*Proof.* We need both  $\beta\alpha = 0$  as well as surjectivity of  $\beta$  and injectivity of  $\alpha$ . The actual equation  $\beta\alpha = 0$  reduces exactly to a quadratic polynomial in the  $w_1, w_2$  with the two ASD equations emerging as coefficients.  $\square$

**Observation 3.5.** *This can be viewed as a complex on the trivial vector bundles  $\underline{V}, \underline{W \oplus V \oplus V}$  over  $\mathbb{C}^2$*

$$\underline{V} \xrightarrow{\alpha} \underline{W \oplus V \oplus V} \xrightarrow{\beta} \underline{V}$$

Now because we have Hermitian structure on each of  $W, V$ , and  $U$ , we have hermitian structure on the space we are interested. We can thus define adjoints  $\alpha^\dagger, \beta^\dagger$ . In particular the Hermitian structure gives us canonical projection operators  $P_\beta$  onto  $\ker \beta$  and  $P_\alpha$  ( $\text{im } \alpha$ ) $^\perp = \ker \alpha$  so that  $P_x = P_{\beta,x} P_{\alpha,x}$  is then a projection onto  $\text{im } \alpha^\perp \cap \ker \beta \cong \ker \beta / \text{im } \alpha$ .

The above proposition also implies

$$\Delta_{x,y}^\dagger := \begin{pmatrix} \beta_{x,y} \\ \alpha_{x,y}^\dagger \end{pmatrix} : W \oplus (V \otimes \mathbb{C}^2) \rightarrow V \times V \quad (26)$$

is a surjection. Explicitly:

$$\Delta_{x,y}^\dagger = \begin{pmatrix} w_2 I + w_1 J^\dagger & w_2 B_2 - w_1 B_1^\dagger + z_2 & w_2 B_1 + w_1 B_2^\dagger + z_1 \\ -\bar{w}_1 I + \bar{w}_2 J^\dagger & -\bar{w}_1 B_2 - \bar{w}_2 B_1^\dagger - \bar{z}_1 & -\bar{w}_1 B_1 + \bar{w}_2 B_2 + \bar{z}_2 \end{pmatrix} \quad (27)$$

Moreover, there is an adjoint operator to  $\Delta^\dagger$  on these bundles:

$$\Delta := (\beta^\dagger \quad \alpha) = \begin{pmatrix} \bar{w}_2 I^\dagger + \bar{w}_1 J & w_2 J - w_1 I^\dagger \\ \bar{w}_2 B_2^\dagger - \bar{w}_1 B_1 + \bar{z}_2 & -w_2 B_1 - w_1 B_2^\dagger - z_1 \\ \bar{w}_2 B_1^\dagger + \bar{w}_1 B_2 + \bar{z}_1 & w_2 B_2 - w_1 B_1^\dagger + z_2 \end{pmatrix} \quad (28)$$

More compactly, if we write

$$a = \begin{pmatrix} I^\dagger & J \\ B_2^\dagger & -B_1 \\ B_1^\dagger & B_2 \end{pmatrix}, \quad b = \begin{pmatrix} 0 & 0 \\ I_k & 0 \\ 0 & I_k \end{pmatrix} \quad (29)$$

then

$$\Delta = aw + bz \quad (30)$$

when we write  $w$  and  $z$  as quaternions in this space by associating to a complex pair  $(q_1, q_2) = q \in \mathbb{C}^2$  the quaternionic operator:

$$q \leftrightarrow \begin{pmatrix} \bar{q}_2 & -q_1 \\ \bar{q}_1 & q_2 \end{pmatrix} \quad (31)$$

for any  $q_1, q_2 \in \mathbb{C}$ . This structure is compatible with the operator  $R$ :

**Proposition 3.6.**  $\Delta_{xq,yq}^\dagger = \bar{q}\Delta_{x,y}^\dagger$

*Proof.* We have that

$$\begin{aligned} \Delta_{x,y}^\dagger &= (awq + bzq)^\dagger \\ &= q^\dagger(aw + bz) \\ &= q^\dagger\Delta^\dagger \end{aligned} \quad (32)$$

□

Define the bundle vector  $E$  at  $(x, y)$  as the vector space corresponding to the kernel of the  $\Delta^\dagger$  map at  $(x, y)$ .

**Corollary 3.7.**  $E_{x,y} = E_{xq,yq}$ , meaning  $x, y$  are projective coordinates over the quaternions.

The above makes  $E$  a bundle on the projective space  $\mathbb{P}^1(\mathbb{H}) \cong S^4$ . On this compact space, we can calculate topological charge.

Because of this symmetry, we can specialize to the case  $y = 1$ , i.e.  $(w_1, w_2) = (0, 1)$  in the ADHM equations. This simplifies the operator  $\Delta^\dagger$  to

$$\Delta^\dagger = \begin{pmatrix} I & B_2 + z_2 & B_1 + z_1 \\ J^\dagger & -\bar{B}_1^\dagger - \bar{z}_1 & \bar{B}_2^\dagger + \bar{z}_2 \end{pmatrix} \quad (33)$$

Solutions to ADHM correspond to  $\Psi$  such that

$$\Delta^\dagger \Psi = 0. \quad (34)$$



It is easy to see that

$$\Delta^\dagger \Delta = \begin{pmatrix} f^{-1} & 0 \\ 0 & f^{-1} \end{pmatrix} \quad (35)$$

for some Hermitian  $f$ . We can also construct an *orthonormal* matrix  $M$  whose columns span  $\ker \Delta^\dagger$ . Clearly then:

$$\Delta^\dagger M = 0.$$

The set of solutions  $\Psi$  to  $\Delta^\dagger \Psi = 0$  gives rise to  $M$  and gives a connection:

$$M^\dagger dM.$$

We can then define the projection operator:

$$Q := \Delta f \Delta^\dagger \quad (36)$$

as well as

$$P := MM^\dagger \quad (37)$$

**Lemma 3.8.**  $P + Q = 1$ . That is,  $P$  projects into the null space of  $\Delta^\dagger$ .

**Proposition 3.9.** This gives rise to a connection  $A = M^\dagger dM$

*Proof.* Take  $s$  a section so that  $Ms$  gives a section on  $E = \ker \Delta^\dagger$ , then

$$\begin{aligned} Mds + MAs &= d_A(Ms) \\ &= Pd(Ms) \\ &= MM^\dagger d(Ms) \\ &= M(ds + (M^\dagger dM)s) \end{aligned} \quad (38)$$

giving our result. □

**Proposition 3.10.**  $A \in \mathfrak{su}(n)$ .

*Proof.*  $A^\dagger = (dM)^\dagger M = -M^\dagger dM$  because of normalization:  $M^\dagger M = 1$ . □

**Proposition 3.11.**  $A$  is anti-self-dual.

*Proof.*

$$\begin{aligned} F_{\mu\nu} &= \partial_{[\mu} A_{\nu]} + A_{[\mu} A_{\nu]} \\ &= \partial_{[\mu} (M^\dagger \partial_{\nu]} M) + (M^\dagger \partial_{[\mu} M)(M^\dagger \partial_{\nu]} M) \\ &= (\partial_{[\mu} M^\dagger)(\partial_{\nu]} M) + (M^\dagger \partial_{[\mu} M)(M^\dagger \partial_{\nu]} M) \\ &= (\partial_{[\mu} M^\dagger)(\partial_{\nu]} M) + (\partial_{[\mu} M^\dagger)M(M^\dagger \partial_{\nu]} M) \\ &= (\partial_{[\mu} M^\dagger)(1 - P)(\partial_{\nu]} M) \\ &= (\partial_{[\mu} M^\dagger)Q(\partial_{\nu]} M) \\ &= (\partial_{[\mu} M^\dagger)\Delta f \Delta^\dagger(\partial_{\nu]} M) \\ &= M^\dagger(\partial_{[\mu} \Delta)f(\partial_{\nu]} \Delta^\dagger)M \end{aligned} \quad (39)$$

The term involving the derivatives of these  $\Delta$  operators

$$(\partial_{[\mu}\Delta)f(\partial_{\nu]}\Delta^\dagger) \quad (40)$$

can be reduced to the action of sigma matrices  $-i\sigma_\mu$  on  $f$ :

$$\begin{aligned} \partial_\mu\Delta &= -i\sigma_\mu \\ \Rightarrow (\partial_{[\mu}\Delta)f(\partial_{\nu]}\Delta^\dagger) &= (-i\sigma_{[\mu} \otimes I_k)(I_2 \otimes f)(-i\sigma_{\nu]}^\dagger \otimes I_k) \\ &= -2i\sigma_{\mu\nu} \otimes f \end{aligned} \quad (41)$$

And we know  $\star\sigma_{\mu\nu} = -\sigma_{\mu\nu}$ . This illustrates how the underlying quaternionic structure gives rise to ASD solutions.  $\square$

**Proposition 3.12.** *The topological charge of  $E$  when considered as a bundle over  $S^4$  is  $-k$*

*Proof.* (Sketch) Note that  $W \oplus (V \otimes U) \cong \mathbb{C}^{n+2k} = E \oplus E^\perp$ . Since  $E$  has dimension  $n$  this leaves a complement of complex dimension  $2k$ . This can be identified as  $k$  one-dimensional copies of the quaternions, so that  $W \oplus (V \otimes U)$  decomposes as a direct sum

$$E \oplus \mathbb{H}^{\oplus k} \quad (42)$$

so corresponds to  $k$  quaternion line bundles over  $S^4$ . In fact this turns out to be the **tautological line bundle**  $\Sigma$ .

Now from simple Chern theory, we know:

$$0 = c_2(\mathbb{C}^{n+2k}) = c_2(E) + kc_2(\Sigma). \quad (43)$$

But the second chern number of a quaternionic tautological bundle is 1 (analogous to how the first chern number of a complex tautological bundle is 1). This gives  $c_2(E) = -k$ .  $\square$

**Corollary 3.13.**  *$A$  is a framed connection, and the topological charge is  $-k$ .*

*Proof.* We know  $A$  over  $\mathbb{R}^4$  extends to a connection over  $S^4 = \mathbb{P}^1(\mathbb{H})$ .  $\square$

## References

- [1] John C Baez and Javier Muniain *Gauge Theory, Knots, and Gravity*: 1994, World Scientific.
- [2] Nick Dorey et al. *The Calculus of Many Instantons*: 2002, hep-th/0206063
- [3] AJ Lindenhovius *Instantons and the ADHM Construction*: 2011
- [4] Alexander Kirillov Jr. *Quiver Varieties and Quiver Representations*: 2016: Graduate Studies in Mathematics, AMS.
- [5] Hiraku Nakajima *Hilbert Schemes of Points on Surfaces* 1991: University Lecture Series, AMS.